

# Zorn's Lemma and the Alexander Subbase Lemma

David Milovich

April 24, 2018

**Lemma 1** (Alexander Subbase Lemma). *Let  $\mathcal{S}$  be a subbase of space  $X$  such that every subcover of  $\mathcal{S}$  has a finite subcover. Then  $X$  is compact.*

*Proof using Zorn's Lemma.* Seeking a contradiction, suppose  $\mathcal{S}$  is as above but  $X$  has an open cover without a finite subcover. Let  $\mathbb{O}$  denote the set of all open covers of  $X$  that lack finite subcovers. Then  $\mathbb{O}$  is nonempty; hence, the union  $\emptyset$  of the empty chain  $\emptyset$  is a subset of some  $\mathcal{U} \in \mathbb{O}$ . By the following claim, each union of a nonempty chain  $\mathbb{C} \subset \mathbb{O}$  is also a subset of some  $\mathcal{U} \in \mathbb{O}$ .

*Claim.* If  $\mathbb{C} \subset \mathbb{O}$  is a nonempty chain, then  $\bigcup \mathbb{C} \in \mathbb{O}$ .

*Proof.* Let  $\mathcal{V} \in \mathbb{C}$  and  $\mathcal{W} = \bigcup \mathbb{C}$ . Then  $\mathcal{W}$  is a set of open subsets of  $X$  because every  $\mathcal{C} \in \mathbb{C}$  is. Moreover,  $\mathcal{W}$  covers  $X$  because  $\mathcal{V}$  does and  $\mathcal{W}$  contains  $\mathcal{V}$ . If  $\{W_1, \dots, W_n\}$  is a finite subcover of  $\mathcal{W}$ , then  $W_i \in \mathcal{C}_i \in \mathbb{C}$  for some  $\mathcal{C}_i$ , for each  $i$ . But in this case, since  $\mathbb{C}$  is a chain,  $\mathcal{C}_1, \dots, \mathcal{C}_n \subset \mathcal{C}_j$  for some  $j$  and, therefore,  $\{W_1, \dots, W_n\}$  is a finite subcover of  $\mathcal{C}_j$ , which is impossible because  $\mathcal{C}_j \in \mathbb{C} \subset \mathbb{O}$ . Therefore,  $\mathcal{W}$  has no finite subcover. Thus,  $\mathcal{W} \in \mathbb{O}$ .  $\square$

By Zorn's Lemma,  $\mathbb{O}$  has a maximal element  $\mathcal{M}$ . Let  $\mathcal{A} = \mathcal{M} \cap \mathcal{S}$ . Since  $\mathcal{A} \subset \mathcal{M} \in \mathbb{O}$ , no finite subset of  $\mathcal{A}$  covers  $X$ . But all subcovers of  $\mathcal{S}$  have finite subcovers; hence,  $\mathcal{A}$  does not cover  $X$ ; choose  $p \in X - \bigcup \mathcal{A}$ . Since  $\mathcal{M}$  does cover  $X$ , we may choose  $M$  such that  $p \in M \in \mathcal{M}$ . Since  $\mathcal{S}$  is a subbase for  $X$ , we may choose  $S_1, \dots, S_n \in \mathcal{S}$  such that  $p \in \bigcap_{i \leq n} S_i \subset M$ .

For each  $S_i$ , we have  $S_i \notin \mathcal{M}$  because  $S_i \in \mathcal{S} - \mathcal{A}$  because  $p \in S_i - \bigcup \mathcal{A}$ . Therefore,  $\mathcal{B}_i = \mathcal{M} \cup \{S_i\} \notin \mathbb{O}$  by maximality of  $\mathcal{M}$ . But since  $\mathcal{B}_i \supset \mathcal{M}$  and every  $S \in \mathcal{S}$  is open in  $X$ , the set  $\mathcal{B}_i$  is still an open cover of  $X$  despite  $\mathcal{B}_i \notin \mathbb{O}$ . Therefore,  $\mathcal{B}_i$  has a finite subcover  $\mathcal{F}_i$ . Let  $\mathcal{G}_i = (\mathcal{F}_i - \{S_i\}) \cup \{M\}$ , which is a finite subset of  $\mathcal{M}$  whose union contains  $Y_i = (X - S_i) \cup M$ .

Let  $\mathcal{G} = \bigcup_{i \leq n} \mathcal{G}_i$ . Then  $\mathcal{G}$  is a finite subset of  $\mathcal{M}$  whose union contains  $\bigcup_{i \leq n} Y_i$ , which, as demonstrated below, is all of  $X$  because  $\bigcap_{i \leq n} S_i \subset M$ .

$$\bigcup_{i \leq n} ((X - S_i) \cup M) = \left( X - \bigcap_{i \leq n} S_i \right) \cup M \supset (X - M) \cup M = X$$

Thus,  $\mathcal{M}$  has a finite subcover despite  $\mathcal{M} \in \mathbb{O}$ . Contradiction!  $\square$

**Lemma 2** (Zorn's Lemma). *Suppose that  $\mathcal{D}$  is set of sets such that, for each chain  $\mathcal{C} \subset \mathcal{D}$ , there exists  $B \in \mathcal{D}$  such that  $\bigcup \mathcal{C} \subset B$ . Then  $\mathcal{D}$  has a maximal element.*

*Proof using the Well-Ordering Principle.* Let  $\leq$  be a well-ordering of  $\mathcal{D}$ . Define  $f: \mathcal{D} \rightarrow \{0, 1\}$  recursively by  $f(B) = 1$  if  $A \subset B$  for all  $A < B$  satisfying  $f(A) = 1$ , and  $f(B) = 0$  otherwise. (If  $f$  were not well-defined, there would be a least  $B$  at which  $f(B)$  was not well-defined. But then  $f(A)$  would be well-defined for all  $A < B$ , implying that our above recursive definition of  $f(B)$  actually does define  $f(B)$ . Therefore,  $f$  must be well-defined.)

For each pair  $A, B \in \mathcal{D}$ , we have  $A \leq B$  or  $A \geq B$ . And if  $A < B$  and  $f(A) = f(B) = 1$ , then  $A \subset B$ . Therefore,  $f^{-1}(\{1\})$  is a chain. Therefore, we may choose  $M \in \mathcal{D}$  such that  $\bigcup (f^{-1}(\{1\})) \subset M$ . Let us show that  $M$  is maximal in  $\mathcal{D}$ . Given  $M \subset C \in \mathcal{D}$ , it is enough to show that  $C \subset M$ . Since  $\bigcup (f^{-1}(\{1\})) \subset M \subset C$ , we have  $f(C) = 1$  by definition of  $f$ . Therefore,  $C \subset \bigcup (f^{-1}(\{1\})) \subset M$ .  $\square$

*Remark.* In the context of  $Z$ , Zermelo's axioms of set theory excluding Choice, the Axiom of the Choice, the Well-Ordering Principle, and Zorn's Lemma are all provably equivalent, but not intuitively equivalent. The old joke is that the Axiom of Choice is obviously true, the Well-Ordering Principle is obviously false, and as for Zorn's Lemma, who can say?